




plain and simple facts

# DRUG INFORMATION



## Anti-HIV Medications (Anti-retroviral Therapy)

Anti-HIV medications are an important part of your treatment against HIV infection. They are medications that slow down the growth of the HIV virus. Because HIV is a special type of virus called “retrovirus,” the medications are usually called “**anti-retrovirals.**”

### What kinds of anti-HIV medications are available?

There are different types (or classes) of anti-HIV medications. Each type of anti-HIV medication works by blocking a different protein that the HIV virus need in its reproduction.

Currently there are two major classes of anti-HIV medications available on the market:

- medications that block the protein “**reverse transcriptase,**” known as **reverse transcriptase inhibitors** or **RT inhibitors (RTIs).**
- medications that block the protein “**protease,**” known as **protease inhibitors (PIs).**

### The RT Inhibitors (nukes and non-nukes)

The RT inhibitors, or RTIs, are divided into 3 groups based on differences in their chemical structures: **nucleoside RT inhibitors** (also called “**nukes**” or **NRTIs**), **non-nucleoside RT inhibitors** (also called “**non-nukes**” or **NNRTIs**), and **nucleotide RT inhibitors** (also called **NtRTIs**).

Medications from the class of **nukes** or **NRTIs** include:

- AZT (also called zidovudine or Retrovir)
- ddI (also called didanosine or Videx or Videx EC)
- ddC (also called zalcitabine or Hivid)
- d4T (also called stavudine or Zerit or Zerit XR)
- 3TC (also called lamivudine or Epivir)
- Abacavir (also called ABC or Ziagen)
- Combivir (a combination medication of AZT + 3TC)
- Trizivir (a combination medication of AZT, 3TC + abacavir)

Medications from the **non-nuke** or **NNRTI** class include:

- nevirapine (also called Viramune)
- delavirdine (also called Rescriptor)
- efavirenz (also called Sustiva)

# Anti-HIV Medications (Anti-retroviral Therapy)

---

The approved medication from the **NtRTI** class is:

- tenofovir (also called Viread)

Medications from the class of **protease inhibitors** include:

- saquinavir (also called Invirase and Fortovase)
- indinavir (also called Crixivan)
- ritonavir (also called Norvir)
- nelfinavir (also called Viracept)
- amprenavir (also called Agenerase)
- Kaletra (lopinavir + ritonavir)

## **When should I start taking anti-HIV medications?**

Starting anti-HIV medications is a big decision. You have to be ready before you start these medications because usually they need to be taken on a regular schedule every day for them to work. This may involve making some changes in your daily routine and lifestyle.

It is important to find a doctor who is knowledgeable about HIV treatments and is sensitive to your situation and culture when working with you in making treatment decisions.

In general, anti-HIV medications are recommended if:

- Your CD4+ cell count (measure of your immune system) is less than 350.
- Your viral load (measure of amount of HIV virus in your blood) is greater than 55,000.
- You have had any symptoms of HIV disease.

## **Which medications should I be taking?**

The current standard of HIV treatment includes the use of a combination of at least 3 different anti-HIV medications from different classes.

There are 2 major reasons for using a combination of anti-HIV medications from different classes together:

- Using different classes of medications enables you to block different proteins that the HIV virus needs at different steps of its reproduction, therefore combination therapy will be more effective than medication from only one class.
- The HIV virus can change itself with every reproduction and may become resistant to certain anti-HIV medications, which means that those medications won't be effective at fighting your virus anymore. When different medications are used together, it is much harder for the virus to change itself to be able to resist all the medications together.

When choosing which medications to be used in your combination, your doctor should work with you to consider a combination that:

- includes different types of anti-HIV medications (that attack the HIV virus at different stages of its reproduction).
- maximizes your ability to take the medications regularly.
- minimizes side effects.
- leaves options open for future changes.

Ask your doctor or pharmacist for detailed information about each of the medications being



recommended for you to take and talk with them about any concerns you have before you start taking these medications.

## **How do I know if the medications are working?**

If your anti-HIV medications are working (that is, if they are effective in controlling the HIV virus), you should expect:

- a decrease in your viral load (ideally to “undetectable” level).
- an increase in your T-cell count.
- fewer infections and symptoms related to HIV.
- better general health.

By going to your doctor regularly to have checkups and blood tests to check your viral load and T-cells, you will get a better idea if the medications are working for you.

## **When do I need to change medications?**

You may need to change the anti-HIV medications in your combination if:

- the medications are not working (this is called “treatment failure”).
- the medications give you very bad side effects.

If you need to change medications because your combination is not working, that is, it’s no longer effective in controlling the virus, your doctor would usually suggest changing at least 2 of the 3 medications in your combination, or maybe all of them. If you have taken many different HIV medications before, your doctor may need to do a special test, called “resistance testing,” before changing your medications. This test lets you and your doctor know which drugs will not work against HIV in your body.

If you are having very bad side effects from your medications, your doctor may suggest decreasing the dose of the medication that is causing the problem or replacing it with another medication. In this case you do not need to change all the medications in your combination because they are still working.

As we learn more about the effects of anti-HIV medications and as newer medications become available, the treatment guidelines may change. Please make sure you work with your doctor and pharmacist to get the most up-to-date information and decide on a medication combination that will work best to meet your specific needs.



---

**Decisions about particular medical treatments should always be made in consultation with a qualified medical practitioner knowledgeable about HIV-related illness and the treatments in question.**



The Canadian AIDS Treatment Information Exchange (CATIE) in good faith provides information resources to help people living with HIV/AIDS who wish to manage their own health care in partnership with their care providers. Information accessed through or published or provided by CATIE, however, is not to be considered medical advice. We do not recommend or advocate particular treatments and we urge users to consult as broad range of sources as possible. We strongly urge readers to consult with a qualified medical practitioner prior to undertaking any decision, use or action of a medical nature.

We do not guarantee the accuracy or completeness of any information accessed through or published or provided by CATIE. Persons relying on this information do so entirely at their own risk. Neither CATIE nor the Public Health Agency of Canada nor any of their employees, directors, officers or volunteers may be held liable for damages of any kind that may result from the use or misuse of any such information. The views expressed herein or in any article or publication accessed or published or provided by CATIE are solely those of the authors and do not reflect the policies or opinions of CATIE or the views of the Public Health Agency of Canada.

Copyright: This Fact Sheet is made available through a collaboration between Asian Community AIDS Services (ACAS) and the Canadian AIDS Treatment Information Exchange (CATIE). Original content developed by ACAS, 2001.

This information is also available in French, Chinese, Tagalog, Vietnamese, Swahili, and other languages at [languages.catie.ca](http://languages.catie.ca) and [www.acas.org](http://www.acas.org)

Funding has been provided by the Public Health Agency of Canada.